

P.O. Box 1646
Castle Rock, Colorado 80104
720-733-0184
Toll Free 1-877-745-3447
info@donatedeggs.com

Donor Number: 0250 (For Agency Use Only)

Today's Date: May 3, 2013

How did you hear of An Eggceptional Match? (If website, pls. specify): Google

Name: _____

Date of Birth: 02/12/1984

Social Security #: _____

Insurance Co: Cigna

Address: _____ City: Glenwood Springs State: CO Zip: _____

Home Phone: _____ Work Phone: _____

Cell Phone: _____ May we leave a Message at (Pls. Circle): Home Work Cell

E-mail Address: _____

I check my email: all day once a day several times a week rarely

Are you currently listed with any other clinics or agencies? Yes If yes, whom? _____

Have you ever been denied entry into another egg donor program? No If yes, please explain in detail:

How soon are you able to begin your donation? ASAP

Who may we contact in case of an emergency? _____ Ph: _____

Who may we contact in case your demographics change? _____ Ph: _____

Are you (Pls. Circle): Married Single with relationship Single without relationship

Are you a U.S. Citizen? Yes No

Do you have medical insurance? Yes Are you willing to travel for an egg donation? Yes

Do you have any legal cases pending against you? No Have you ever filed bankruptcy? Yes

Have you ever been convicted of a crime? No If yes, please elaborate: _____

PHYSICAL CHARACTERISTICS

Age: 29 Height: 5'6" Weight: 120 lbs **Measurements:** Bust 36 Hips 33? Waist 30?

Race: Caucasian (Caucasian, African American, Asian, Hispanic, etc...)

Ethnicity: (Pls. Be specific, ie...German, French, Irish, etc...) _____

Mother's Side: English

Father's Side: Irish

Blood Type: A (+ or -)

Place of Birth: KC, MO

What celebrity do people most commonly say you look like? Reese Witherspoon

Please circle appropriate response:

Body Type/Bone Structure: small **medium** large

Hands: **right-handed** left-handed ambidextrous

| | | | | | |
|-------|--------|--------------|----------------|---------------|-------------|
| Eyes: | *Color | brown | hazel | green | blue |
| | *Set | narrow | average | wide | |
| | *Size | small | average | large | |
| | *Shape | round | oval | almond | |
| | *Shade | light | medium | dark | |

| | | | | | | |
|-------|-----------------|-----------------|---------------|--------------|-----|-------------|
| Hair: | *Natural Color | blond | brown | black | red | other _____ |
| | *Color as child | blond | brown | black | red | |
| | *Shade | light | medium | dark | | |
| | *Type | straight | wavy | curly | | |
| | *Fullness | thin | medium | thick | | |
| | *Texture | fine | medium | course | | |

| | | | | |
|-------|----------------|---------------|----------------|-------|
| Nose: | *Size | small | medium | large |
| | *Width | narrow | average | wide |
| | *length | short | average | wide |
| | *Nostril Flare | small | average | wide |

| | | | | |
|-------------|-------------|--------|----------------|--------|
| Cheekbones: | *Set | low | average | high |
| | *Prominence | slight | medium | strong |

| | | | | |
|--------|-------|-------|----------------|-------------|
| Mouth: | *Size | small | average | large |
| | *Lips | thin | average | full |

| | | | | |
|-------|-------------|-------------|----------------|--------|
| Chin: | *Shape | square | oval | round |
| | *Prominence | slight | average | strong |
| | *Cleft | none | slight | medium |

| | | | | | |
|------------------------|---|----------|-------------|---------------|-------------|
| Other Facial Features: | *Moles | none | one | several | numerous |
| | *Freckles | none | several | moderate | numerous |
| | *Dimples | none | slight | medium | deep |
| Eyesight: | *Vision | normal | far-sighted | near-sighted | |
| | *Glasses | none | single | bifocal | |
| | *Astigmatism | yes | no | age diagnosed | <u>21</u> |
| Dental: | *Device | none | braces | retainer | other _____ |
| | *Reason | cosmetic | accident | disease | other _____ |
| | *Age during use _____ to _____ years of age | | | | |
| Other: | *List _____ | | | | |
| | *Reason/Cause _____ | | | | |

Age at first period? 13 Are your cycle's regular? Yes

How long are your cycles from day one to the next day one? 30-31 How long do they last? 5 Days

Do you experience cramps? None **Mild** Average Severe

Method of birth control? None If none, in the past? _____

Have you ever been pregnant? No If yes, did you have trouble conceiving? _____

Have you ever been treated for infertility? No

Did your mother take DES while she was pregnant with you? No

| Year | Delivery ♀ or ♂ Section/Vag | Miscarriage | Ectopic | Blighted Ovum | Termination |
|------|--------------------------------|-------------|---------|---------------|-------------|
| 1. | | | | | |
| 2. | | | | | |
| 3. | | | | | |
| 4. | | | | | |

Any complications? _____

MEDICAL HISTORY

Have you ever had any surgeries? If so please list type and date:

1. Tonsils, Adenoids (1994)
2. L5-S1 Disc Herniation-(2003)

Have you had a blood transfusion in the last 12 months? No

If yes, please list date and reason: _____

Any hospitalizations not mentioned above? No If yes, please explain: _____

Have you been exposed to radiation or toxic chemicals in your work or personal life? No

Have you received a bite from an animal suspect for rabies within the last 6 months? No

Have you ever had a reaction to anesthesia? No If yes, please explain reaction in detail: _____

*Do you smoke cigarettes? No Packs per day? _____ # of years _____ # of years quit _____

Do you now or have you ever taken recreational drugs? No If so, What? _____

Do you drink alcohol? Yes If yes, how many drinks per: day? _____ week? _____ month? 1

Do you have any allergies to drugs or environmental exposures? No Pls. explain: _____

Describe any childhood allergies that you have outgrown: None

Do you have any medical illnesses (diabetes, asthma, etc...)? N/A If yes, pls. explain: _____

Please list all prescription or over the counter medications including dosage you are currently taking: N/A

***To become an egg donor, you must be completely drug free. You must also be nicotine free for at least 6 mos. You may be tested for this substance at any time.**

Have you ever donated your eggs before? Yes If yes, Please list dates and outcomes:

| Mo/Year | # Eggs Retrieved | # Eggs Fertilized | Did a pregnancy occur? | Did a live birth occur? |
|---------|------------------|-------------------|------------------------|-------------------------|
| 06/2008 | 16 | 11 | Yes-Twins | Yes (Twin Boys) |
| 11/2009 | 20 (18 mature) | 14 | Yes-Singleton | Yes (Boy) |
| 2/2010 | 20 | 14 | Yes-Singleton | Yes |
| 10/2010 | 19 | 19 | Yes | Yes |
| | | | | |

Were their embryos left to cryopreserve (freeze)? Yes If yes, approximately how many per cycle: Cycle #2- 7 embryos
Cycle #3-5 embryos

What is the compensation you are asking for your donation? \$10,000 (1st time donors \$5,000)

What is the least amount you would consider? \$10,000

Will you require missed wages from work? Yes

If yes, what is your hourly wage? \$18.00

How many hours per week do you work? 50+

Will you require childcare reimbursement? No

If yes, what is the hourly rate? _____ X _____ kids

Have you been sexually active in the past 6 months? Yes

Are you currently sexually active? Yes If yes, is it a monogamous relationship and for how long? 6 years
If no, will your partner consent to standard blood testing? _____

Have you or your partner ever had a sexually transmitted disease? No If yes, when and what was your treatment regimen? _____

Have you ever been diagnosed with any gynecological problems such as endometriosis, ovarian cysts, abnormal Pap smears, fibroids, polyps? No If yes, please explain treatment _____

Please mark any that apply to you within the last 12 months:

- ☐ Exposure to HIV
- ☐ Exposure to Hepatitis B or C
- ☐ Had sex in exchange for money or drugs
- ☐ Intravenous drug use
- ☐ Piercing or tattoos
- ☒ None of the Above

EDUCATION

Highest Level of Education Completed: Grade School _____
Jr. High School _____
Sr. High School (GPA: 3.6)
Currently in College pursuing a degree in: _____
Completed College with degree in: A.S. Business Administration
Currently pursuing advanced degree in: Fed. Acq. And Contract Management
Completed advance degree in: _____
Vocational/Trade School: _____

Test Scores: SAT's: _____ ACT's: 29 College GPA: 3.4

Please list names and year of all colleges attended:

| <u>College</u> | <u>Year</u> |
|-------------------|--------------------|
| 1. <u>Park</u> | <u>2003-2006</u> |
| 2. <u>Strayer</u> | <u>2006-2011</u> |
| 3. <u>GCU</u> | <u>2013-Presnt</u> |

What was your favorite subject in school? English/History You're least favorite? Geometry

Dean's List or Honor Roll? Yes

As an adult I am most proud of: School, career, Moves from KC to DC to CO

Currently I have a career in: Business Consulting

I have been in this profession for 3 days/mos/years

*I have flexibility in my current profession: Yes No

Languages: Speak: _____
Read: _____
Write: _____

I consider myself: Athletic Active Average Inactive

Physical activities include: Tennis, horseback riding and jogging

Have you excelled in any physical activities? Yes

Manual Dexterity: Dexterous Average Clumsy

I would describe my diet as: Good

Other skills or talents? _____

Do you show artistic or musical ability? Yes If yes please explain: Violin, Piano, Viola, Clarinet and Flute

***Once you are chosen as an egg donor, first morning appointments are mandatory. You are not able to choose the dates of your appointments, as they will coincide with the clinic's requirements and timing of your cycle. You will be required to attend 7-10 monitoring appointments prior to the actual egg retrieval. Most clinics are open between 7:30-8:00 a.m. If your schedule cannot accommodate this criterion, you will not qualify for the egg donation process.**

FAMILY HEALTH HISTORY

| | Hair Color | Eye Color | Height Weight | Skin Tone | Age If Living | Age at Death | Cause of Death |
|--------------------------|---------------|--------------|------------------|--------------|------------------|-----------------|--|
| Mother | Brown | Brown | 5'5" 140 | Med | 54 | | |
| Father | Red | Hazel | 6'0" 200 | Fair | 58 | | |
| Brother: 1. | Red | Hazel | 5'8" 145 | Fair | 17 | | |
| 2. | | | | | | | |
| 3. | | | | | | | |
| 4. | | | | | | | |
| Sister: 1. | | | | | | | |
| 2. | | | | | | | |
| 3. | | | | | | | |
| 4. | | | | | | | |
| Maternal Grandmother | Lt. Brown | Hazel | 5'7" 140 | Med | | 34 | Uterine Infection-Lived in country. Dr. did not make house call in time. |
| Maternal Grandfather | Brown | Blue | 6'0" 215 | Med | 87 | | |
| Paternal Grandmother | Black | Brown | 5'7" 145 | Med | 85 | | |
| Paternal Grandfather | Red | Blue | 6'0" 215 | Med | | 84 | Heart Attack |
| Children: (If Any) 1. | | | | | | | |
| 2. | | | | | | | |
| 3. | | | | | | | |
| 4. | | | | | | | |

Are you adopted? No If yes, do you have access to your biological health history? _____

Twins or multiple births in the family? No If yes, how many sets? _____

Are there any known genetic diseases that run in your family? No If yes, please identify all such diseases and explain in as much detail as possible:

Has anyone in your family been born with a birth defect? No If yes, please explain in detail: _____

Have you had a brother or sister die in infancy or early childhood? No If yes, please explain the cause of death:

Have you ever been tested for:

Cystic Fibrosis (Caucasian) Not Sure

Sickle Cell (African American) _____

Thalassemia (Greek/Italian) _____

Tay-Sach's (Jewish) _____

If yes to any of the above, were you determined a carrier? _____

How would you describe your personality and temperament? Easy going, positive.

What is your philosophy of life? Take it as it comes....You determine the outcome by your own actions.

What qualities and characteristics would you hope the recipient parents possess? Supportive, patience, understanding and intelligent.

How does it make you feel at the possibility of their offspring knowing about the donation? Proud that I was able to help!

How would you describe your childhood? Great! Awesome, supportive parents!

What is the earliest memory you hold as a child? Going to the lake with the entire family.

What was it like growing up in your family? Wonderful! Great family.

What religion did you belong to as a child? Catholic

When I Was A Child:

My favorite thing to do was: Play with my cat

At home I was expected to: Clean up after myself, behave.

My parents were strict about: Bedtime and school

My parents taught me to value: Life!

What I loved most about my father was: Sense of humor

What I loved most about my mother was: Her ability to always listen

My favorite relatives were: Gramms, P-Aunt

I loved to visit: Aunt Judith

In comparison to others I was: Well off

Your Teenage Years:

Describe yourself as a teenager: Adventurous

Describe your achievements: Good grades, a lot of true friendships

Did you do poorly at anything? No

Did you have any problems as a teenager (e.g., health, acne, social, educational, etc.)? No

What do you hope to achieve by volunteering in an egg donor program? (e.g., emotionally, financially): Helping someone who truly needs help....financially, just to invest in my future.

What message would you like passed on to the recipient of your eggs/their offspring? How eager I was to help and how happy I am for them!

What helped you decide to become an egg donor? Rewarding and the effect it had on me the first time

Do you consider yourself a reliable person? Yes

Do you consider yourself a punctual person? Yes

Would you describe yourself as a religious or spiritual person? Somewhat

Do you have any ethical, moral or religious reservations about being an egg donor? No

What are your personal goals? Have you achieved any of these goals? Advance in my career and education. So far I have reached #3 on my list of 10! Right on track....

What do you see yourself doing in the next 5-10 years? Advancing my career and having a family.

What would you like your recipient couple to know about you that has not already been asked? Nothing I can think of...

What is your favorite color? Green

Favorite type of food? Mom's!

Favorite movie? Annie

Favorite type of music? I like them all!

Favorite Book? The Thin Woman-Dorothy Cannell

Would you be willing to donate to gay or single prospective parents? Yes Please specify: Race, color, creed nor sexuality matter to me.

Would you be willing to meet a child conceived as the result of your donation? Yes Please elaborate: If that was their wish, absolutely!

Would you be interested in possibly meeting the prospective parents? If they would like!

What are your beliefs concerning selective reduction or aborting a fetus due to an anomaly(s) (birth defect)?

I believe if it is for better for all involved, it should be done.

How do you feel about the possibility of any remaining embryos being donated to another infertile couple that cannot afford the cost of infertility treatment? Will you sign a consent permitting such donation?

I am requesting to share excess eggs in an effort to preserve my future fertility. Up for discussion on details of this arrangement.

How do you feel about any remaining embryos being discarded or donated to scientific research? Will you sign a consent permitting such medical or scientific research?

Please see above.

Some clinics have their Prospective Parents sign away rights to any left over embryos to be used at the clinic's discretion, how do you feel about not knowing the outcome of their decision?

Please see above.

What are your thoughts on the possibility of this couple using a gestational carrier (surrogate) or a qualified sperm donor?

If it helps them achieve their dream, I don't mind.

Is there a message you would like to leave for your prospective parents? Best of luck!!

Please read the list of medical problems carefully and indicate your answer in the appropriate box. Remember, this is the only information that your intended parents have to make a healthy and intuitive choice for their offspring.

| HEART | You | Mother | Father | Siblings | Grandparents | Other Family | Explain |
|--|-----|--------|--------|----------|--------------|--------------|---------|
| A. Stroke | | | | | | | |
| B. heart attack | | | | | | | |
| C. heart disease | | | | | | | |
| 1. from birth | | | | | | | |
| 2. lifestyle | | | | | | | |
| D. hardening of the arteries | | | | | | | |
| E. high blood pressure | | | | | | | |
| BLOOD | | | | | | | |
| A. anemia | | | | | | | |
| B. sickle-cell anemia | | | | | | | |
| C. hemophilia or other bleeding problem | | | | | | | |
| D. leukemia | | | | | | | |
| E. Immune Deficiency | | | | | | | |
| F. other blood disorder | | | | | | | |
| RESPIRATORY (LUNGS) | | | | | | | |
| A. hay fever | | | | | | | |
| B. asthma | | | | | | | |
| C. emphysema | | | | | | | |
| D. tuberculosis | | | | | | | |
| E. lung cancer | | | | | | | |
| F. pneumonia | | | | | | | |
| G. other lung disease | | | | | | | |
| GASTRO-INTESTINAL | | | | | | | |
| A. ulcer of stomach or duodenum | | | | | | | |
| B. gall stones | | | | | | | |
| C. hepatitis A | | | | | | | |
| D. hepatitis B | | | | | | | |
| E. cirrhosis | | | | | | | |
| F. colon cancer | | | | | | | |
| G. ulcerative colitis | | | | | | | |
| H. Crohn’s disease | | | | | | | |
| I. cystic fibrosis | | | | | | | |
| J. intestinal cancer | | | | | | | |
| K. any other cancer/digestive prob. | | | | | | | |
| METABOLIC/ENDOCRINE | | | | | | | |
| A. diabetes mellitus | | | | | | | |
| B. hypoglycemia | | | | | | | |
| C. thyroid cancer | | | | | | | |
| D. thyroid disease | | | | | | | |
| E. goiter | | | | | | | |
| F. adrenal dysfunction or disorder | | | | | | | |
| G. hyperactivity | | | | | | | |
| URINARY | | | | | | | |
| A. kidney disease | | | | | | | |
| B. other disease of urinary tract (urethra, bladder, ureter) | | | | | | | |
| GENITAL/REPRODUCTIVE | | | | | | | |
| A. undescended testicle | | | | | | | |
| B. hypospadias | | | | | | | |
| C. prostate cancer | | | | | | | |
| D. uterine fibroids | | | | | | | |
| E. ovarian cysts | | | | | | | |
| F. cancer of cervix, ovaries or uterus | | | | | | | |

| | You | Mother | Father | Siblings | Grandparents | Other Family | Explain |
|--|-----|--------|--------|----------|--------------|--------------|---------|
| NEUROLOGICAL | | | | | | | |
| A. migraines | | X | | | | | |
| B. mental retardation | | | | | | | |
| C. senility before age 50 | | | | | | | |
| | | | | | | | |
| D. Multiple Sclerosis | | | | | | | |
| E. Cerebral Palsy | | | | | | | |
| F. epilepsy | | | | | | | |
| G. hydrocephalus | | | | | | | |
| H. disorder of the spinal cord | | | | | | | |
| I. Huntington's chorea | | | | | | | |
| J. Gaucher's disease | | | | | | | |
| K. Wilson's disease | | | | | | | |
| L. Creutzfeldt-Jacob disease | | | | | | | |
| M. Alzheimer's disease | | | | | | | |
| N. other diseases of the nervous system | | | | | | | |
| MENTAL HEALTH | | | | | | | |
| A. schizophrenia | | | | | | | |
| B. bipolar or manic depressive | | | | | | | |
| C. depression | | | | | | | |
| MUSCLE/BONE/JOINTS | | | | | | | |
| A. muscular dystrophy | | | | | | | |
| B. other chronic muscle disease | | | | | | | |
| C. lupus | | | | | | | |
| D. deformity of the spine | | | | | | | |
| E. osteoporosis | | | | | | | |
| F. dwarfism | | | | | | | |
| G. heredity low back disease | | | | | | | |
| H. arthritis | | | | | | | |
| I. gout | | | | | | | |
| SIGHT/SOUND/SMELL | | | | | | | |
| A. deafness before age 60 | | | | | | | |
| B. deformity of the ear | | | | | | | |
| C. cataracts before age 50 | | | | | | | |
| D. blindness | | | | | | | |
| E. color blindness | | | | | | | |
| F. glaucoma | | | | | | | |
| G. deviated septum | | | | | | | |
| H. any other sight/sound/smell disorders | | | | | | | |
| SKIN | | | | | | | |
| A. acne | | | | | | | |
| B. eczema | | | | | | | |
| C. skin cancer | | | | | | | |
| D. pigmentation disorders | | | | | | | |
| E. other disorders of the skin | | | | | | | |
| OTHER | | | | | | | |
| A. alcoholism | | | | | | | |
| B. drug abuse, misuse or addiction | | | | | | | |
| C. breast cancer | | | | | | | |
| D. any other cancer not mentioned above | | | | | | | |
| E. any other condition not mentioned above | | | | | | | |

No To all

| RISK FACTORS | Yes | No | Comment |
|---|------------|-----------|----------------|
| Have you ever been sexually active with a male who was gay or bisexual? | Yes | No | |
| Have you ever injected drugs or had a sexual partner who did so? | Yes | No | |
| Have you ever had hemophilia or received any human derived clotting factor concentrates, including factor VIII or factor IX concentrate? | Yes | No | |
| Have you ever had a sexual partner with hemophilia or who received any human derived clotting factor concentrates? | Yes | No | |
| Have you ever had sex in exchange for money or drugs? | Yes | No | |
| Have you ever been sexually active with a person who has had sex in exchange for money or drugs? | Yes | No | |
| Have you ever been sexually active with a person who was known or suspected to have HIV, hepatitis B or hepatitis C? | Yes | No | |
| Have you been exposed to body fluids, open wounds, non-intact skin or mucus membranes of any person known or suspected to have HIV, hepatitis B and/or C? | Yes | No | |
| Have you had an accidental needle stick within the past 12 months? | Yes | No | |
| Have you ever been or have you had a sexual partner who was incarcerated for 72 consecutive hours or longer? | Yes | No | |
| In the past 12 months, have you lived with or had contact with anyone known or suspected to have hepatitis? | Yes | No | |

(Cont'd)

| | | |
|---|------------|-----------|
| Have you acquired a tattoo or other skin piercing procedure within the preceding 12 months? | Yes | No |
|---|------------|-----------|

| | | |
|--|------------|-----------|
| Have you ever been diagnosed with hepatitis? | Yes | No |
|--|------------|-----------|

| | | |
|---|------------|-----------|
| Have you been vaccinated or had contact with anyone vaccinated for smallpox within the past 2 months? | Yes | No |
|---|------------|-----------|

| | | | |
|---|------------|-----------|---------------------|
| Have you ever been diagnosed with or suspected to have West Nile Virus? | Yes | No | if so, when? |
|---|------------|-----------|---------------------|

| | | | |
|--|------------|-----------|---------------------|
| Have you ever been treated for or diagnosed with chlamydia, gonorrhea, herpes or syphilis? | Yes | No | if so, when? |
|--|------------|-----------|---------------------|

| | | | |
|---|------------|-----------|--------------------|
| Have you or any of your blood relatives been diagnosed and/or have a history of transmissible spongiform encephalopathy such as Creutzfeldt-Jakob disease or variant Creutzfeldt-Jakob disease? | Yes | No | if so, who? |
|---|------------|-----------|--------------------|

| | | |
|---|------------|-----------|
| Have you ever received a non-synthetic dura mater transplant or a pituitary-derived growth hormone? | Yes | No |
|---|------------|-----------|

| | | |
|--|------------|-----------|
| Do you have a history of changes in cognition, speech or gait? | Yes | No |
|--|------------|-----------|

| | | | |
|---|------------|-----------|----------------------|
| Have you ever received a blood transfusion? | Yes | No | if so, where? |
|---|------------|-----------|----------------------|

| | | |
|--|------------|-----------|
| Have you visited or lived in the United Kingdom for three months or more between 1980-1996 including England, Scotland, Wales, Ireland, Isle of Man, Channel Islands, Gibraltar or Falkland Islands? | Yes | No |
|--|------------|-----------|

| | | |
|--|------------|-----------|
| Were you a member of the US military, civilian military, employee or a dependent of a member of the military stationed in Belgium, the Netherlands, Germany, Spain, Portugal, Turkey, Italy or Greece between 1980-1996? | Yes | No |
|--|------------|-----------|

(Cont'd)

| | | | |
|--|------------|-----------|----------------------|
| From 1980 to present, have you spent time that adds up to 5 years or more in Europe? | Yes | No | if so, where? |
|--|------------|-----------|----------------------|

| | | | |
|---|------------|-----------|---------------------|
| Were you born in or have you lived in any of the following Countries since 1977; Cameroon, Central Africa Republic, Chad, Congo, Equatorial, Guinea, Gabon, Niger or Nigeria? | Yes | No | If so, when? |
|---|------------|-----------|---------------------|

| | | |
|--|------------|-----------|
| If yes, were you given a blood transfusion or any medical treatment with a product made from blood while you were there? | Yes | No |
|--|------------|-----------|

| | | |
|---|------------|-----------|
| Have you ever had sexual contact with anyone who was born or lived in Cameroon, Central Africa Republic, Chad, Congo, Equatorial, Guinea, Gabon, Niger or Nigeria since 1977? | Yes | No |
|---|------------|-----------|

| | | | |
|--|------------|-----------|---------------------|
| Have you or someone you know been diagnosed, treated or suspected of having sudden acute respiratory syndrome? (SARS)? | Yes | No | if so, when? |
|--|------------|-----------|---------------------|

| | | | |
|--|------------|-----------|--------------------|
| Have you, your sexual partner, and/or anyone you live with ever had a transplant or other medical procedure that involves being exposed to live cells, tissues or organs from an animal? | Yes | No | if so, who? |
|--|------------|-----------|--------------------|

| | | |
|--|------------|-----------|
| Have you been exposed to blood, saliva or fluids from the person described in the proceeding question? | Yes | No |
|--|------------|-----------|

| | | |
|---|------------|-----------|
| Have you ever received a human organ, tissue transplant or human extract? | Yes | No |
|---|------------|-----------|

| | | | |
|---|------------|-----------|--------------------|
| Have you ever been excluded as a blood donor? | Yes | No | if so, why? |
|---|------------|-----------|--------------------|

| | | |
|---|------------|-----------|
| Have you been diagnosed or suspected to have Chagas' disease? | Yes | No |
|---|------------|-----------|

(Cont'd)

Have you been exposed to significant levels of radiation, toxic chemicals, or heavy metals (such as lead, mercury or gold) in your home or work environment?

Yes No

Have you received a bite from an animal suspected for rabies within the last six months?

Yes No

CONSENT

Under the penalty of perjury, I attest that all of the information I have provided in my donor application is true and complete to the best of my knowledge. I also have a good understanding of the commitment involved in the egg donation process.

Donor's Printed Name: _____

Donor's Signature: _____

Date: _____

I _____ give An Eggceptional Match, LLC full authority to include my photographs on its web site, and to share my personal profile with prospective parents but without disclosing to them any of my identifying information. I hereby release An Eggceptional Match, LLC and Angela Bevill, and their agents, employees, successors and assigns, from any and all liability of any nature whatsoever as a result of such disclosures.

Donor's Signature: _____

Date: _____

Witness to Signatures above: _____

Date: _____

DEFINITIONS

Inherited – A disease or characteristic that is transmitted through genes from parents to offspring. Inheritance patterns include the following:

Autosomal Dominant – Disorders caused by one mutated copy of a gene. An affected person usually has one affected parent. Autosomal dominant disorders usually occur in every generation of an affected family. When a person carries an autosomal dominant gene mutation, each of his/her offspring has a 50% chance for inheriting the gene mutation.

Autosomal Recessive – Disorders caused by two mutated copies of a gene. An affected person usually has unaffected parents who each carry one copy of the mutated gene. Autosomal recessive disorders are not usually seen in every generation of a family. Carrier parents have a 25% chance for having an affected child.

X-linked dominant – Disorders caused by mutations in genes located on the X chromosome. Females are more frequently affected than males, and the chance to pass on an X-linked dominant disorder differs between men and women. Fathers cannot pass the X-linked traits or disorders to their sons. Females who have an X-linked dominant gene mutation have a 50% chance to have an affected child.

X-linked recessive – Disorders caused by mutations on genes on the X chromosomes. Males are more often affected than females, and the chance to pass on the disorder differs between men and women. Families with X-linked recessive disorders often have affected males, but rarely affected females, in each generation. Females who carry an X-linked recessive gene mutation have a 50% chance to pass it on to each of her children.

Multifactorial – Disorders caused by a combination of the effects of multiple genes or by interactions between genes and the environment.

Sources and additional information:

Talking Glossary of Genetic Terms <http://www.genome.gov/10002096>; <http://www.genome.gov/glossary.cfm#g>

Fact Sheets <http://www.genome.gov/10000202>

Cancer Dictionary <http://www.cancer.gov/dictionary/>

Genetics Home Reference National Library of Medicine <http://ghr.nlm.nih.gov/>

National Institutes of Health Genetic and Rare Diseases Information Center

<http://rarediseases.info.nih.gov/GARD/Default.aspx?PageID=4>

Gene Tests <http://www.genetests.org/>

Cancer

- **Hereditary Breast/Ovarian Cancer** – Mutations in *BRCA1* or *BRCA2* genes predispose to breast cancer and ovarian cancer as well as prostate cancer (*BRCA1*) and other cancers (*BRCA2*). Hereditary breast/ovarian cancer is inherited in families in an autosomal dominant pattern. Each child of an individual with a *BRCA1* or *BRCA2* cancer-predisposing mutation has a 50% chance of inheriting the mutation.
- **Hereditary colon cancer**
 - **Hereditary non-polyposis colorectal cancer** - Hereditary non-polyposis colon cancer (HNPCC) is caused by an autosomal dominant inherited gene mutation. HNPCC is characterized by an increased risk of colon cancer and other cancers (e.g., of the endometrium, ovary, stomach, small intestine, hepatobiliary tract, upper urinary tract, brain, skin). Each child of an individual with a HNPCC cancer-predisposing mutation has a 50% chance of inheriting the mutation.
 - **Heart**
- **Congenital heart disease** - Congenital heart disease is a common type of birth defect or malformation in one or more structures of the heart or blood vessels that occurs during pregnancy while the fetus is developing. The cause of congenital heart disease is not known in most affected people. There are some recognized factors that are associated with an increased risk for congenital heart disease including: 1) genetic or chromosomal abnormalities such as Down syndrome; 2) taking certain medications, alcohol or drug abuse during pregnancy; and 3) maternal viral infections such as German measles in the first trimester of pregnancy. The risk of having a child with congenital heart disease is higher if a parent or a sibling has a congenital heart defect.

Blood

- **Sickle cell anemia** - Sickle cell disease is a group of disorders that affects hemoglobin, the molecule in red blood cells that delivers oxygen to cells throughout the body. Individuals who have sickle cell disease have atypical hemoglobin molecules called hemoglobin S, which can distort red blood cells into a sickle, or crescent, shape. Signs and symptoms include a low number of red blood cells (anemia), repeated infections, and periodic episodes of pain. The severity of symptoms varies from person to person. Sickle cell anemia is inherited in an autosomal recessive manner. Each child of carrier parents has a 25% chance to be born with sickle cell anemia.
- **Factor V Leiden thrombophilia** - Factor V Leiden thrombophilia is an inherited disorder of blood clotting. Factor V Leiden is the name of a specific mutation that results in thrombophilia - the increased tendency to form abnormal blood clots in blood vessels. People who have the factor V Leiden mutation are at somewhat higher than average risk for a type of clot that forms in veins, such as the deep veins of the legs (deep venous thrombosis), or a clot that travels through the bloodstream and lodges in the lungs (pulmonary embolism). Factor V Leiden thrombophilia can be inherited in families in an autosomal dominant and autosomal recessive manner.
- **Hemophilia** - Hemophilia is a bleeding disorder that slows the blood clotting process. People who have hemophilia often experience prolonged bleeding or oozing following an injury, surgery, or having a tooth pulled. The major types of this condition are hemophilia A (also known as classic hemophilia) and hemophilia B (also known as Christmas disease). Hemophilia A and hemophilia B are inherited in an X-linked recessive manner. In X-linked recessive inheritance, a female with one altered copy of the gene in each cell is called a carrier. She can pass on the altered gene to her children, but usually does not experience signs and symptoms of the disorder.
- **Tay-Sachs** - Tay-Sachs disease is a rare inherited disorder that causes progressive destruction of nerve cells in central nervous system (the brain and spinal cord). Affected infants progressively lose motor skills such as turning over, sitting, and crawling. Children who have the severe infantile form of Tay-Sachs disease usually survive only into early childhood. Tay-Sachs disease is inherited in an autosomal recessive manner. Carrier parents have a 25% in each pregnancy to have an affected child.
- **Thalassemia** - Beta thalassemia is an inherited blood disorder that reduces the production of hemoglobin. Symptoms of beta thalassemia occur when not enough oxygen gets to various parts of the body due to low levels of hemoglobin and a shortage of red blood cells. Beta thalassemia is inherited in an autosomal recessive manner. Carrier parents have a 25% chance in each pregnancy to have an affected child.

Respiratory

- **Alpha-1 antitrypsin disorder** - Alpha-1 antitrypsin deficiency is an inherited condition that can cause lung disease in adults and liver disease in adults and children. This disorder is inherited in an autosomal co-dominant pattern. Co-dominance means that two different versions of the gene may be expressed, and both versions contribute to the genetic trait.

Gastrointestinal

- **Cystic Fibrosis** - Cystic fibrosis is an inherited disorder of the mucus glands that affects many body systems. The most common signs and symptoms of cystic fibrosis include progressive damage to the respiratory system and chronic digestive system problems. Cystic fibrosis is inherited in an autosomal recessive manner. Carrier parents have a 25% chance in each pregnancy for having an affected child.
- **Pyloric stenosis** - Pyloric stenosis (also called infantile pyloric stenosis or gastric outlet obstruction) is a condition that involves a narrowing of the pylorus, the lower part of the stomach through which food and other stomach contents pass to enter the small intestine. When an infant has pyloric stenosis, the muscles in the pylorus become enlarged to the point where food is prevented from emptying out of the stomach. Pyloric stenosis is known to run in families. When a parent has pyloric stenosis, then, their infant has an increased risk of developing the disorder.

Metabolic/Endocrine

- **Phenylketonuria** - Phenylketonuria (also known as PKU) is an inherited disorder that increases the levels of a substance called phenylalanine in the blood. Phenylalanine is a building block of proteins that is obtained through the diet. If PKU is not treated, phenylalanine can build up to harmful levels in the body, causing mental retardation and other serious health problems. PKU is inherited in an autosomal recessive manner. Carrier parents have a 25% chance with each pregnancy to have an affected child.
- **Dwarfism** – There are a number of different types of dwarfism and many are inherited in families. Examples of types of dwarfism include: achondroplasia, thanatophoric dysplasia, and Robinow syndrome.

Urinary

- **Polycystic kidney disease** - Polycystic kidney disease is a disorder that affects the kidneys and other organs. Cysts, develop in the kidneys, causing them to become enlarged and can lead to kidney failure. Cysts may also develop in other organs, particularly the liver. There are two major forms of polycystic kidney disease distinguished by the age of onset and their pattern of inheritance. The autosomal dominant form (sometimes called ADPKD) has signs and symptoms that typically begin in adulthood, although cysts in the kidney are often present from childhood. The autosomal recessive form of polycystic kidney disease (sometimes called ARPKD) is much rarer and is often lethal early in life.

Genital/Reproductive

- **Hypospadias** – Hypospadias is a birth defect of the urethra that happens in males. It involves an abnormally placed opening in the penis. Instead of opening at the tip of the penis, a hypospadiac urethra opens anywhere along the line running from the tip along the underside of the shaft to the where the penis and scrotum meet. In most males
- hypospadias is not inherited, nor is their family recurrence. In some cases, hypospadias happens as a result of a chromosomal abnormality called a pericentric inversion of chromosome number 16.

Reproductive Outcomes

- **2 or more miscarriages** – Miscarriage (also called spontaneous abortion) is the term used for a pregnancy that ends on its own, within the first 20 weeks of gestation. The causes of miscarriages are varied, and most often the cause cannot be identified. During the first trimester, the most common cause of miscarriage is chromosomal abnormality - meaning that something is not correct with the baby's chromosomes. In some cases the chromosomal abnormality in the developing fetus is the result of a parent carrying a balanced chromosomal arrangement called a translocation. This can lead to multiple miscarriages.

- **Birth defects** – A birth defect is a problem that happens while the baby is developing in the mother's body. Most birth defects happen during the first 3 months of pregnancy. A birth defect can affect almost any part of the body. Causes of birth defects include a family history of birth defects, maternal age, certain drugs taken during pregnancy, alcohol use and smoking during pregnancy.

Neurological

- **Mental Retardation** - Mental retardation is a term used to describe a person who has certain limitations in mental functioning and difficulties in communicating, taking care of him or herself, and social skills. These limitations will cause a child to learn and develop more slowly than a typical child. Causes of mental retardation include genetic conditions such as Down syndrome, problems during pregnancy, problems at birth and health problems such as malnutrition.
- **Cerebral palsy** - Cerebral palsy is the term for a group of disorders that involve the loss of movement or loss of other nerve function. Cerebral palsy is caused by injuries to the largest part of the brain (cerebrum) which happen as the baby grows in the womb or near the time of birth. There are multiple causes of cerebral palsy including birth defects that affect the brain, spinal cord, head, face, lungs or metabolism, and certain hereditary and genetic conditions.
- **Neurofibromatosis** – There are two types of neurofibromatosis. Neurofibromatosis type 1 is a disorder characterized by changes in skin coloring (pigmentation) and the growth of tumors along nerves in the skin, brain, and other parts of the body. The signs and symptoms of this condition vary widely among affected people. Neurofibromatosis type 1 is considered to have an autosomal dominant pattern of inheritance. Neurofibromatosis type 2 is a disorder characterized by the growth of noncancerous tumors in the nervous system. Neurofibromatosis type 2 is also considered to have an autosomal dominant pattern of inheritance. However, unlike most other autosomal dominant conditions, in which one altered copy of a gene in each cell is sufficient to cause the disorder, two copies of the NF2 gene must be altered to trigger tumor formation in neurofibromatosis type 2. A mutation in the second copy of the NF2 gene happens in other cells in the nervous system during a person's lifetime. Almost everyone who is born with one NF2 mutation acquires a second mutation in many cells and develops the tumors characteristic of neurofibromatosis type 2.
- **Autism/Aspergers** –
 - Autism and autism spectrum disorders are complex neurodevelopmental conditions. The genetics of autism are complex and it is thought that there are multiple genes involved.
 - Aspergers – Asperger syndrome is one of several autism spectrum disorders, with symptoms of difficulty in social interactions and restricted, stereotyped interests and activities. Children who have Aspergers syndrome do not usually have language or cognitive developmental delays. Genes are believed to play a role in Aspergers syndrome, and it seems to run in some families.
- **Hydrocephalus** – Hydrocephalus is a condition in which the primary characteristic is excessive accumulation of fluid in the brain. The excessive accumulation of fluid causes an abnormal widening of spaces in the brain called ventricles. This widening creates potentially harmful pressure on the tissues of the brain. The causes of hydrocephalus are still not well understood. Hydrocephalus may be caused by inherited genetic abnormalities (such as the genetic defect that causes aqueductal stenosis) or developmental disorders (such as those associated with neural tube defects including spina bifida and encephalocele). Other possible causes include complications of premature birth, and diseases such as tumors or hemorrhage which block the fluid.
- **Tuberous sclerosis** - Tuberous sclerosis is a genetic disorder characterized by the growth of numerous noncancerous tumors in many parts of the body. These tumors can occur in the skin, brain, kidneys, and other organs, in some cases leading to significant medical problems. Tuberous sclerosis is inherited in an autosomal dominant manner, which means one copy of the altered gene in each cell is sufficient to cause the disorder. In about one-third of families, an affected person inherits an altered gene from a parent who has the disorder. About two thirds of cases result from new gene mutations. These cases occur in people with no history of tuberous sclerosis in their family.
- **Creutzfeldt-Jakob Disease** – Creutzfeldt-Jakob disease is a prion disease. Prion diseases are group of progressive conditions that affect the nervous system. Prion diseases impair brain function, causing memory changes, personality changes, a decline in intellectual function, and problems with movement that worsen over time. The signs and symptoms of these conditions usually begin in adulthood, and these disorders lead to death within a few months to several years. Only a small percentage of prion disease cases run in families. Most cases occur in people without any known risk factors or gene mutations. Creutzfeldt-Jakob disease is acquired by eating beef products obtained from cattle that have prion disease.

- **Huntington Disease** - Huntington disease is a progressive brain disorder that causes uncontrolled movements, mental and emotional problems, and loss of thinking ability. Adult-onset Huntington disease, is the most common form of this disorder, with onset usually in a person's thirties or forties. An early-onset, less common form of Huntington disease begins in childhood or adolescence. This condition is inherited in an autosomal dominant manner, which means one copy of the altered gene in each cell is sufficient to cause the disorder.
- **Gaucher Disease** - Gaucher disease is an inherited disorder that affects many of the body's organs and tissues. The signs and symptoms of this condition vary widely among affected individuals. There are several types of Gaucher disease based on their particular features. Some types do not affect the brain and spinal cord while others do. Type 1 Gaucher disease, for example, is the most common form of this disorder. Major signs and symptoms of Type 1 Gaucher disease include enlargement of the liver and spleen, a low number of red blood cells, easy bruising caused by a decrease in blood platelets, lung disease, and bone abnormalities such as bone pain, fractures, and arthritis. Types 2 and 3 Gaucher disease, on the other hand, have problems that affect the central nervous system. Gaucher disease is inherited in an autosomal recessive pattern, which means both copies of the gene in each cell have mutations. The parents each carry one copy of the mutated gene, but they do not show signs or symptoms of the disease.
- **Wilson's Disease** - Wilson disease is an inherited disorder in which excessive amounts of copper accumulate in the body, particularly in the liver, brain, and eyes. Typically, signs and symptoms of Wilson disease first appear during the teenage years. Wilson's disease is inherited in an autosomal recessive pattern, which means both copies of the gene in each cell have mutations. The parents each carry one copy of the mutated gene, but they do not show signs or symptoms of the disease.
- **Tourette syndrome** - Tourette syndrome is a complex disorder characterized by repetitive, sudden, and involuntary movements or noises called tics. Tics usually appear in childhood, and their severity varies over time. In most cases, tics become milder and less frequent in late adolescence and adulthood. Individuals who have Tourette syndrome are also at risk for other associated problems including attention deficit hyperactivity disorder (ADHD), obsessive-compulsive disorder (OCD), anxiety, depression, and problems with sleep. A variety of genetic and environmental factors appear to play a role in causing Tourette syndrome. Most of these factors are unknown to date. Among family members of an affected person, it is therefore difficult to predict who else may be at risk of developing the condition.

Mental Health

- **Depression** – Clinical depression is an illness that can challenge a person's ability to perform even routine daily activities, and in some cases lead a person to contemplate or commit suicide. There are several different types of depression (mood disorders that include depressive symptoms) such as major depression, bipolar disorder and seasonal depression. The causes of depression are complex. Genetic, biological, and environmental factors can contribute to its development. In some people, depression can be traced to a single cause, while in others, a number of causes are involved. For many, the causes are never known. Certain types of depression seem to run in some families. Research is ongoing as to exactly which genes are involved in depression.

Muscle/Bone Joint

- **Muscular dystrophy** - Muscular dystrophies are a group of genetic conditions characterized by progressive muscle weakness and wasting. The Duchenne and Becker types of muscular dystrophy primarily affect the skeletal muscles, which are used for movement, and the muscles of the heart. These conditions occur much more frequently in males than in females. Both Duchenne and Becker muscular dystrophy are inherited in an X-linked recessive pattern, with the mutated gene that causes the disorder on the X chromosome. Males are affected by X-linked recessive disorders much more frequently than females.
- **Achondroplasia** - Achondroplasia is a disorder of bone growth, particularly in the long bones of the arms and legs. All people with achondroplasia have short stature. Health problems commonly associated with achondroplasia include breathing difficulties (called apnea), obesity, and recurrent ear infections. Achondroplasia is inherited in an autosomal dominant pattern, which means one copy of the altered gene in each cell is sufficient to cause the disorder. About 80 percent of individuals with achondroplasia have average-size parents; these cases result from a new gene mutation in that individual. In the remaining cases, people with achondroplasia have inherited a gene from one or two affected parents.

- **Osteogenesis imperfecta** - Osteogenesis imperfecta (OI) is a group of genetic disorders that mainly affect the bones. People who have OI have bones that break easily, often from mild trauma or with no apparent cause. Multiple fractures are common, and in severe cases, fractures can occur even before birth. Milder cases may involve only a few fractures over a person's lifetime. There are at least eight recognized forms of osteogenesis imperfecta, designated type I through type VIII, distinguished by their signs and symptoms. Most types of osteogenesis imperfecta have an autosomal dominant pattern of inheritance, which means one copy of the altered gene in each cell is sufficient to cause the disorder. Many people with type I or type IV osteogenesis imperfecta inherit a mutation from a parent who has the disorder.
- **Marfan syndrome** - Marfan syndrome is a connective tissue disorder. Connective tissue provides strength and flexibility to structures throughout the body such as bones, ligaments, muscles, the walls of blood vessels, and heart valves. Marfan syndrome affects most organs and tissues, especially the skeleton, lungs, eyes, heart, and the large blood vessel that distributes blood from the heart to the rest of the body called the aorta. Individuals who have Marfan syndrome often are tall and slender, have elongated fingers and toes, a long narrow face, highly arched palate, and have an arm span that exceeds their body height. About half of all people with Marfan syndrome have vision problems caused by a dislocated lens (ectopia lentis). Most people with Marfan syndrome have abnormalities of the heart and the aorta. This condition is inherited in an autosomal dominant pattern, which means one copy of the altered gene in each cell is needed to cause the disorder. At least one quarter of classic Marfan syndrome cases result from a new gene mutation. These individuals have no history of the disorder in their family.
- **Spinal muscular atrophy** - Spinal muscular atrophy is a disorder that affects the control of muscle movement. It is caused by a loss of specialized nerve cells, (motor neurons), in the spinal cord and the part of the brain that is connected to the spinal cord (the brainstem). The loss of motor neurons leads to weakness and shrinkage of muscles used for activities such as crawling, walking, sitting up, and controlling head movement. In severe cases of spinal muscular atrophy, the muscles used for breathing and swallowing are affected. There are a number of different subtypes of spinal muscular atrophy based on the age of onset and symptoms. Most types of spinal muscular atrophy are inherited in an autosomal recessive pattern, which means both copies of the gene in each cell have mutations. One type of spinal muscular atrophy is inherited in an autosomal dominant manner, which means one copy of the altered gene in each cell is sufficient to cause the disorder.
- **Reiter's disease** - Reiter's syndrome, also known as reactive arthritis, is a type of arthritis that occurs as a reaction to an infection somewhere in the body. Most infections that cause the disease begin in the bladder, urethra, penis, or vagina and are spread through sexual intercourse, a form of the disease called genitourinary Reiter's syndrome, or urogenital Reiter's syndrome. Other infections that can cause reactive arthritis include gastrointestinal infections due to eating contaminated food or handling contaminated substances, a form of the disease called gastrointestinal Reiter's syndrome, or enteric Reiter's syndrome. Reiter's syndrome affects mostly young men, between the ages of 20 and 40. Although researchers are not sure why some people develop reactive arthritis in response to certain infections, a genetic factor (presence of the HLA-B27 gene) appears to increase the risk.

Sight/Sound/Smell

- **Deafness** – There are several types of deafness including conductive hearing loss, neural hearing loss (nerve deafness), and mixed hearing loss (a combination of conductive and neural hearing loss). Some people are born deaf. Usually the cause is unknown. Although deafness is inherited in some families, deaf parents often have hearing children and hearing parents often have deaf children. Diseases and injuries of the ear can also cause deafness.
- **Blindness** – Blindness is a condition of lacking visual perception that is due to physiological or neurological factors. Blindness has a number of causes including disease and malnutrition. Blindness may have a genetic cause, and may also be a symptom of a particular genetic disorder. Recent advances in mapping of the human genome have identified genetic causes of low vision or blindness, for example the disorder called Bardet-Biedl syndrome.
- **Color blindness** – Color blindness is the inability to perceive differences between some of the colors that other people can distinguish. It is usually genetic in nature, but may also be due to eye, nerve or brain damage, or to exposure to certain chemicals. Color blindness can be inherited in families. Since the mapping of the human genome there have been many causative gene mutations discovered. Mutations capable of causing color blindness originate from at least 19 different chromosomes and many different genes.

- **Retinoblastoma** - Retinoblastoma is a rare type of eye cancer that develops in the retina, the part of the eye that detects light and color. Although this disorder can occur at any age, it usually develops in young children. Most cases of retinoblastoma occur in only one eye, but both eyes can be affected. Retinoblastoma can be inherited in an autosomal dominant pattern which means that one copy of the altered gene in each cell is sufficient to increase cancer risk. A person with retinoblastoma may inherit an altered copy of the gene from one parent, or the altered gene may be the result of a new mutation. For retinoblastoma to develop, a second mutation in the other copy of the RB1 gene must occur in retinal cells during the person's lifetime. When there is a family history of retinoblastoma or if the person develops tumors in both eyes, the gene mutation is probably in all of the person's cells, and that person is said to have an inherited form of retinoblastoma. A smaller number of individuals have retinoblastoma as a result of missing portions of chromosome 13 that are not inherited.

Skin

- **Albinism** – Albinism is a condition in which there is a lack of melanin pigment in the eyes, skin and hair (or more rarely the eyes alone). Albinism is hereditary and results from inheritance of recessive gene mutations. There are two main categories of Albinism - 1) **oculocutaneous albinism** in which there is a lack of melanin pigment in skin and hair, and 2) **ocular albinism**, in which only the eyes lack pigment. People with oculocutaneous albinism can have anywhere from no pigment at all to almost-normal levels. People who have ocular albinism have generally normal skin and hair color, and many even have a normal eye appearance. Albinism may also be a feature of a genetic syndrome such as Hermansky-Pudlak syndrome.
- **Neurofibromatosis** – There are two types of Neurofibromatosis – Type 1 and Type 2. Neurofibromatosis type 1 is a disorder characterized by changes in skin coloring and the growth of tumors along nerves in the skin, brain, and other parts of the body. The signs and symptoms of this condition vary widely among affected people. Neurofibromatosis type 2 is a disorder characterized by the growth of noncancerous tumors in the nervous system. The most common develop along the nerve that carries information from the inner ear to the brain (the auditory nerve). Tumors that occur on nerves in other areas of the brain or spinal cord are also commonly seen with this condition. Both Type 1 and Type 2 Neurofibromatosis are considered to have an autosomal dominant pattern of inheritance. People with Neurofibromatosis Type 1 and Type 2 are born with one mutated copy of either the NF1 or NF2 mutated genes in each cell. In about half of cases, the gene mutation is inherited from an affected parent. The remaining cases result from new mutations in the gene and occur in people with no history of the disorder in their family. Unlike most other autosomal dominant conditions, in which one altered copy of a gene in each cell is sufficient to cause the disorder, two copies of either the NF1 or NF2 gene must be altered to trigger tumor formation in neurofibromatosis. A mutation in the second copy of the NF1 or NF2 gene occurs during a person's lifetime in specialized cells surrounding nerves. Almost everyone who is born with one NF1 or NF2 mutation acquires a second mutation in many cells and develops the tumors characteristic of the disease.

Congenital Abnormalities/Birth Defects

- **Cleft lip/palate** – Cleft lip and palate are common birth defects that affect the upper lip and the roof of the mouth. There are many causes of cleft lip and palate. Gene alterations passed down from one or both parents, drugs used or maternal viruses during pregnancy can cause cleft lip and/or palate. Cleft lip and palate can also be part of a genetic syndrome or occur with other birth defects. Risk factors for cleft lip and palate also include a family history of cleft lip or palate and other birth defects.
- **Congenital hip problems** – Congenital hip problems, also called hip dysplasia, involve problems with formation of the hip joint in children. The location of the hip dysplasia can be either the ball of the hip joint (femoral head), the socket of the hip joint (the acetabulum), or both. Hip dysplasia, called congenital dysplasia of the hip (or CDH) in the past is now called developmental dysplasia of the hip (DDH). There are a number of factors that contribute to cause DDH. One known risk factor is having a family history of hip dysplasia. Other causes include when the baby is born in breech position or when there is a lack of intrauterine fluid (oligohydramnios) during pregnancy.

- **Club feet** – Clubfoot is a condition where the foot turns inward and downward. It is a congenital condition, meaning it is present at birth. Other terms for clubfoot are Talipes equinovarus and Talipes. Clubfoot is the most common congenital disorder involving the legs, and can range from mild and flexible to severe and rigid. Although the exact cause is not known, clubfoot may be passed down in some families. Family history, therefore, is a risk factor for clubfoot, as is being a male.
- **Heart Defect** - A congenital heart defect involves an abnormal structure of the heart that is present at birth. Congenital heart defects are the most common type of major birth defect. There are multiple causes of congenital heart defects including environmental and genetic factors. Genes that can cause congenital heart defects are now being discovered, such as a gene that can cause an atrial septal defect and one that may contribute to hypoplastic left heart syndrome. Congenital heart defects can also be a part of a wider pattern of birth defects and genetic syndromes such as Down syndrome, Turner syndrome and velocardiofacial syndrome.
- **Hearing problems** - There are several types of hearing loss including conductive hearing loss, neural hearing loss (nerve deafness), and mixed hearing loss (a combination of conductive and neural hearing loss). Some people are born with hearing loss. Usually the cause is unknown. Although hearing loss is inherited in some families, deaf parents often have hearing children and hearing parents often have deaf children. Diseases and injuries of the ear can also cause deafness.
- **Spina bifida** - Spina bifida is one of a group of birth defects called neural tube defects. Spina bifida occurs during fetal development when a portion of the neural tube fails to develop or close properly causing defects in the spinal cord and in the bones of the backbone. Spina bifida, like many other birth defects appears to be caused by a combination of genetic and environmental risk factors, such as a family history of neural tube defects, folic acid deficiency, and medical conditions such as diabetes and obesity.
- **Microcephaly** - Microcephaly is disorder in which the circumference of the head is smaller than normal because the brain has not developed properly or has stopped growing. Microcephaly can be present at birth or it may develop in the first few years of life. It is most often caused by genetic abnormalities that interfere with the growth of the cerebral cortex during the early months of fetal development. Microcephaly is associated with genetic syndromes such as Down syndrome, chromosomal syndromes, and neurometabolic syndromes, Babies may also be born with microcephaly if their mother abuses drugs or alcohol during pregnancy, or becomes infected with the German measles, chicken pox.
- **Holoprosencephaly** - Holoprosencephaly is a disorder caused by the failure of the embryonic forebrain (*prosencephalon*) to divide properly into the double lobes of the cerebral hemispheres. As a result, the baby has a single-lobed brain structure and severe skull and facial defects. In most cases of holoprosencephaly, the malformations are so severe that babies die before birth. In less severe cases, babies are born with normal or near-normal brain development and facial deformities that may affect the eyes, nose, and upper lip. Often, no specific cause for holoprosencephaly can be identified. There are some specific chromosomal abnormalities that have been identified as the cause of holoprosencephaly in some patients. In some families, holoprosencephaly is inherited in autosomal dominant or X-linked recessive inheritance. Several genes have also been identified that play a role in causing holoprosencephaly.

Chromosomal Abnormalities

- **Down syndrome** - Down syndrome is a chromosomal disorder that is associated with mental retardation, a characteristic facial appearance, and poor muscle tone in infancy. Individuals who have Down syndrome may also have heart defects, digestive problems such as gastroesophageal reflux or celiac disease, hearing loss, and cancer of blood-forming tissue (leukemia). Some people with Down syndrome have hypothyroidism. Down syndrome also appears to be associated with an increased risk of Alzheimer disease. Down syndrome is usually caused by the presence of an extra chromosome number 21, called trisomy 21, which means each cell in the body has three copies of chromosome 21 instead of the usual two copies. Most cases of Down syndrome are not inherited, but occur as random events during the formation of egg or sperm. One type of Down syndrome, called translocation Down syndrome, can be inherited.

- **Fragile X syndrome** - Fragile X syndrome is a genetic disorder that involves a range of developmental problems including learning disabilities and mental retardation, and behavioral problems such as hyperactive behavior and attention deficit disorder. Males are usually more severely affected by this disorder than females. Many males with fragile X syndrome have characteristic physical features that become more apparent with age such as a long and narrow face, large ears, prominent jaw and forehead, unusually flexible fingers, and enlarged testicles after puberty. Most cases of fragile X syndrome are caused by a mutation in which a DNA segment, known as the CGG triplet repeat, is expanded within the FMR1 gene. Fragile X syndrome is inherited in families in an X-linked dominant pattern.
- **Turner syndrome** - Turner syndrome is a chromosomal disorder that affects development in females. Women with Turner syndrome are often shorter than average and are usually unable to conceive children because they lack ovarian function. Other features of Turner syndrome can include extra skin on the neck, puffiness or swelling of the hands and feet, skeletal abnormalities, heart defects, and kidney problems. Developmental delays, learning disabilities, and behavioral problems may also be present, although these characteristics vary among affected females. In most cases, Turner syndrome is not inherited. Rather, it occurs as random events during the formation of egg or sperm.
- **Klinefelter syndrome** - Klinefelter syndrome is a chromosomal disorder that affects male sexual development. Most males who have Klinefelter syndrome have one extra copy of the X chromosome in each cell. The presence of an extra X chromosome interferes with male sexual development causing their testicles to develop abnormally, and leading to low levels of the hormone testosterone beginning during puberty. A lack of testosterone can lead to breast development, reduced facial and body hair, and an inability to father children. Boys who have Klinefelter syndrome may have learning disabilities and difficulty with speech and language development. Klinefelter syndrome is caused by the presence of one or more extra copies of the X chromosome in a male's cells. Klinefelter syndrome is not inherited, but usually occurs as a random event during the formation of egg or sperm.

Genetic History

- **Bloom syndrome** - Bloom syndrome is an inherited disorder that is characterized by a high frequency of breaks and rearrangements in an affected person's chromosomes. Individuals who have Bloom syndrome are usually much smaller than average, and often have a high-pitched voice and characteristic facial features including a long, narrow face; small lower jaw; and prominent nose and ears. They tend to develop pigmentation changes that often appear as a butterfly-shaped patch of reddened skin on the face. Other features of the Bloom syndrome may include learning disabilities, mental retardation, chronic lung problems, diabetes, and immune deficiency that leads to recurrent pneumonia and ear infections. Men with Bloom syndrome are usually not able to father children because they do not produce sperm. Women with the disorder generally experience menopause earlier than usual. Chromosome instability in Bloom syndrome also results in a high risk of cancer in affected individuals. Bloom syndrome is inherited in families in an autosomal recessive pattern, which means both copies of the gene in each cell have mutations.
- **Canavan disease** - Canavan disease is an inherited disorder that causes progressive damage to nerve cells in the brain. The signs and symptoms of Canavan disease usually begin in early infancy; however, the course of the disorder can be quite variable. Infants with Canavan disease usually appear normal for the first few months of life. By age 3 to 5 months, these infants begin to have developmental delays in motor skills, weak muscle tone, large head size, and mental retardation. They may also develop feeding and swallowing difficulties, seizures, and sleep disturbances. Canavan disease is inherited in an autosomal recessive pattern, which means both copies of the gene in each cell have mutations.
- **Fabry Disease** - Fabry disease is an inherited disorder that begins in childhood and results from the buildup of a particular type of fat in the body's cells. Characteristic features of Fabry disease include episodes of pain, particularly in the hands and feet; clusters of small, dark red spots on the skin; a decreased ability to sweat; cloudiness of the front part of the eye; and hearing loss. Individuals with Fabry disease are also at risk for potentially life-threatening complications such as progressive kidney damage, heart attack, and stroke. Fabry disease is inherited in an X-linked pattern; however, unlike other X-linked disorders, Fabry disease causes significant medical problems in many females who have one altered copy of the mutated gene. These women may experience many of the classic features of the disorder.
- **Familial Dysautonomia** - Familial dysautonomia is a genetic disorder that affects the development and survival of certain nerve cells. The disorder causes disturbances in autonomic nerve cells, which control involuntary actions such as digestion, breathing, production of tears, and the regulation of blood pressure and body temperature. It also affects activities related to the senses, such as taste and the perception of pain, heat, and cold. Familial dysautonomia is also called hereditary sensory and autonomic neuropathy, type III. Problems related to this disorder first appear during infancy and include poor muscle tone, feeding difficulties, poor growth, lack of tears, frequent lung infections, and

difficulty maintaining body temperature. Developmental delays in walking and speech, are usually present, although some affected individuals do not show signs of developmental delay. Familial dysautonomia is inherited in an autosomal recessive pattern, which means both copies of the gene in each cell have mutations.

- **Familial Mediterranean Fever** - Familial Mediterranean fever is an inherited disorder that involves recurrent episodes of painful inflammation in the abdomen, chest, or joints. These episodes are often accompanied by fever and sometimes a rash. The first episode usually occurs by the age of 20. For some affected individuals, however, the initial episode occurs much later in life. The episodes usually last 12 to 72 hours and may vary in severity and length of time between attacks. A buildup of protein deposits occurs in some cases of familial Mediterranean fever and this can lead to kidney failure if left untreated. This condition is inherited in an autosomal recessive pattern, which means both copies of the gene in each cell have the mutations. Rarely, familial Mediterranean fever may be inherited in an autosomal dominant pattern, which means one copy of an altered gene is sufficient to cause the disorder.
- **Fanconi Anemia** - Fanconi anemia is a rare, inherited blood disorder that causes bone marrow failure. Fanconi anemia causes the bone marrow to stop making enough new blood cells for the body to function normally. Infants born with Fanconi anemia are at higher risk for having birth defects. Fanconi anemia can also cause the bone marrow to make many abnormal blood cells, which can lead to serious health problems such as cancer. This condition is inherited in an autosomal recessive pattern, which means both copies of the gene in each cell have the mutations.
- **Niemann-Pick, Type A** - Niemann-Pick disease is an inherited disorder that involves lipid metabolism - the breakdown, transport, and use of fats and cholesterol in the body. In affected individuals the abnormal lipid metabolism causes harmful amounts of lipids to accumulate in the spleen, liver, lungs, bone marrow, and brain. There are four main types of Niemann-Pick disease. Type A presents during infancy and is characterized by an enlarged liver and spleen, failure to thrive, and progressive deterioration of the nervous system. Children born with Niemann-Pick, Type A generally do not survive past early childhood. Niemann-Pick, Type A is inherited in an autosomal recessive pattern, which means both copies of the gene in each cell have mutations.
- **Mucopolysaccharidosis Type IV** - Mucopolysaccharidosis Type IV is a genetic disorder, primarily which is characterized by severe neurological and ophthalmologic abnormalities. Also known as ML4, the disorder usually presents during the first year of life with mental retardation, corneal opacities, and delayed motor milestones. Children with ML4 begin to show signs of developmental delay during their first year of life. They usually attain a maximum developmental age of 15 months in language and motor function, although their receptive abilities are more advanced. They may also experience retinal degeneration that severely limits their vision. ML4 is inherited in an autosomal recessive pattern which means both copies of the gene in each cell have mutations.